



barchart(g10 ~ x | g2, subset = g10 != "1")




barchart(unclass(Titanic))


0 200 400 600 800




barchart(g2)




barchart(VADeaths)




barchart(x)




barchart(UCBAdmissions)



bwplot(g2 ~ x + y, outer = TRUE)




bwplot(y, notch = TRUE)







densityplot(~x, groups = g2)



densityplot(y, plot.points = "jitter")




dotplot(g10 ~ x | g2)




dotplot(unclass(Titanic))


0 200 400 600




dotplot(g2)




dotplot(VADeaths)



dotplot(x)




dotplot(UCBAdmissions)







histogram( $\sim c(x, y)$ )




histogram(g2)




histogram(c(x, y))




qqmath( $\sim x + y$ )




qqmath(x)




stripplot(g2 ~ x + y, outer = TRUE)



stripplot(y, jitter = TRUE)




qq(g2 ~ x)







xyplot(y ~ x)



xyplot(ts(x))




levelplot(y ~ g2 + g10)




# levelplot(UCBAdmissions)


Admitted Rejected




levelplot(as.table(x), ...)



levelplot(VADeaths)




contourplot(y ~ g2 + g10)



contourplot(UCBAdmissions)

Admitted Rejected



Female

Male

Gender

Female

Male

Admitted

Rejected


Admitted

Rejected


Admit




contourplot(unclass(Titanic))




contourplot(VADeaths)




cloud( $g_{10} \sim x + y$ )




cloud(VADeaths)




cloud(UCBAdmissions)




wireframe(y ~ g2 + g10)



wireframe(VADeaths)




splom(~cbind(x = x, y = y, g = as.numeric(g2)))



Scatter Plot Matrix




splom(cbind(x = x, y = y, g = as.numeric(g2)))




Scatter Plot Matrix

splom(data.frame(x, y, g2))




Scatter Plot Matrix


parallelplot(~iris)



parallelplot(data.matrix(iris[1:4]))



parallelplot(iris)



rfs(oneway(y ~ g2))

0.2

0.4

0.6

0.8

Fitted Values minus Mean

Residuals

0.2

0.0

-0.2

-0.4


0.2

0.4


0.6

0.8


f-value



tmd(sort(y) ~ sort(x))




tmd(xyplot(sort(y) ~ sort(x)))







xyplot(y ~ x, pch = 16, cex = 1.5)



xyplot(y ~ x | g2, data = g, cex = c(1, 2))



densityplot(g\$x, plot.points = FALSE)



plot(equal.count(rnorm(1000)))

