



barchart(g10 ~ x | g2, subset = g10 != "1")




barchart(unclass(Titanic))


0 200 400 600 800




barchart(g2)



barchart(VADeaths)



barchart(x)



barchart(UCBAdmissions)

0 500 1000 1500

Male


Female

Rejected


Admitted

0 500 1000 1500


Freq




bwplot(g2 ~ x + y, outer = TRUE)




bwplot(y, notch = TRUE)




densityplot(~x, groups = g2)



densityplot(y, plot.points = "jitter")




dotplot(g10 ~ x | g2)




dotplot(unclass(Titanic))


0 200 400 600




dotplot(g2)




dotplot(VADeaths)




dotplot(x)




dotplot(UCBAdmissions)




histogram($\sim c(x, y)$)




histogram(g2)




histogram(c(x, y))




qqmath($\sim x + y$)




qqmath(x)




stripplot(g2 ~ x + y, outer = TRUE)




stripplot(y, jitter = TRUE)




qq(g2 ~ x)




xyplot(y ~ x)



xyplot(ts(x))




levelplot(y ~ g2 + g10)




levelplot(UCBAdmissions)


Admitted Rejected



levelplot(VADeaths)




contourplot(y ~ g2 + g10)



contourplot(UCBAdmissions)

Admitted Rejected



Female

Male

D

E

F

A

B

C

Female

Male

Admitted


Rejected

Admitted


Rejected

Admit


contourplot(unclass(Titanic))




contourplot(VADeaths)




cloud($g_{10} \sim x + y$)




cloud(VADeaths)




cloud(UCBAdmissions)




wireframe(y ~ g2 + g10)



wireframe(VADeaths)




splom(~cbind(x = x, y = y, g = as.numeric(g2)))




Scatter Plot Matrix

splom(cbind(x = x, y = y, g = as.numeric(g2)))




Scatter Plot Matrix

splom(data.frame(x, y, g2))




Scatter Plot Matrix


parallelplot(~iris)



parallelplot(data.matrix(iris[1:4]))



parallelplot(iris)



rfs(oneway(y ~ g2))

0.2


0.4

0.6


0.8

Fitted Values minus Mean


Residuals




tmd(sort(y) ~ sort(x))



tmd(xyplot(sort(y) ~ sort(x)))



xyplot(y ~ x, pch = 16, cex = 1.5)



xyplot(y ~ x | g2, data = g, cex = c(1, 2))

0.0 0.2 0.4 0.6 0.8

1

2

y

x

0.8

0.6

0.4

0.2


0.0

0.2


0.4

0.6

0.8



densityplot(g\$x, plot.points = FALSE)



plot(equal.count(rnorm(1000)))

